Index

```
A_{+}(x) The set of states reachable from
          x by CM(F), 154
A_{\perp}^{k}(x) The set of states reachable from
          x at time k by CM(F), 154
C_{r_0}^k Generalized controllability matrix.,
          159
*, Convolution operator, 76
A \rightsquigarrow B, Uniformly accessibility, 93
A \stackrel{a}{\leadsto} B B is uniformly accessible using
          a from A, 123
A \otimes B, 409
C^{\infty}, Functions whose derivatives of ar-
          bitrary order exist, and are con-
          tinuous, 29
C_V(r) := \{x : V(x) \le r\} The rth sub-
          level set of V, 194
C_n, Controllability matrix, 98
Co, Complex plane, 144
F_k, The output maps for the linear con-
          trol model, 30
G_C^{(r)}(x,B) = \mathsf{E}_x \Big[ \sum_{k=0}^{\sigma_C} 1\!\!1_B(\Phi_k) r^k \Big], 372
I_B, 73
I_{a}(x,B), 133
K_a(x,A) := \sum_{n=0}^{\infty} P^n(x,A)a(n), 123
K_{a_{\varepsilon}}, Resolvent kernel, 69
L(x, A) := P_x(\tau_A < \infty), 71
L(x,h), 297
M_n(g) Martingale derived from g, 439
N(t), Number of customers in a queue
          at time t, 46
N(t), Number of customers in a queue
```

immediately before the nth ar-

ter the n^{th} service time is com-

 N_n^* , Number in a queue immediately af-

rival, 47

pleted, 48

 O_w Control set., 156, 160

```
O_w, Supports disturbance or control in
           CM(F) and NSS(F) models,
           33
P(x,A), n step transition probability,
P(x,A), One step transition probabil-
           ity, 59, 67
P(z) := \sum_{n=0}^{\infty} p(n)z^n, 182
P^n(x,A), n step transition probability,
P_h(x,A), Kernel for "process on h",
Q(x, A), P_x \{ \Phi \in A \text{ i.o.} \}, 204
Q(x,h), 297
R(x), Conditional emptying time for a
           dam model, 52
R_n Residual service time immediately
           after a customer arrival, 78
Re^n, n-dimensional Euclidean space, 6
S_n(g) Partial sum of g(\Phi_k), 415
T(x,A), continuous component, 131
T_{ab} = \min\{j : Z_a(j) = Z_b(j) = 1\},\ 321
U(x, A) := \sum_{n=1}^{\infty} P^{n}(x, A), 71
U(z) := \sum_{n=0}^{\infty} u(n)z^{n}, 182
U_{A}(x, B) := \sum_{n=1}^{\infty} {}_{A}P^{n}(x, B), 74
U_{C}^{(r)}(x, B) = \mathsf{E}_{x} \Big[ \sum_{k=1}^{\tau_{C}} \mathbb{1}_{B}(\Phi_{k})r^{k} \Big], 369
U_{\alpha}^{(r)}(x,f) := \mathsf{E}_x \Big[ \sum_{n=1}^{\tau_{\alpha}} f(\Phi_n) r^n \Big], 361
U_h, Resolvent kernel, 295
V^+(n), Forward recurrence time chain,
V^+(t) := \inf(Z_n - t : Z_n > t, \ n \ge 1)
           Forward recurrence time pro-
           cess, 77
V_{\delta}^{+}(n) = V^{+}(n\delta), 77
V^{-}(t) := \inf(t - Z_n : Z_n \le t, \ n \ge 1)
           Backward recurrence time pro-
```

cess, 77

 $V_{\delta}^{-}(n) = V^{-}(n\delta), 77$

 V_C , Minimal solution to (V2), 271

 $Z_n := \sum_{i=0}^n Y_i$ Delayed renewal process, 76

 $\Delta V(x) = \int P(x, dy)V(y) - V(x)$ Drift operator, 178

 Δ_k Derivative process, 170

 Γ , The distribution of a disturbance variable W, 24

 $\mathcal{G}^+(\gamma)$: distributions which have Laplace-Stieltjes transform convergent in $[0,\gamma],\ 394$

 $\Lambda_i^*(x, A) = P(i, x; 0, A)$ Ladder chain transition probabilities, 78

 $\Lambda_{i-j+1}(x, A) = P(i, x; j, A)$ Ladder chain transition probabilities, 78

 $\Omega = \mathsf{X}^{\infty}$, Sequence space, 56

 $\Omega_+(C)$ Omega limit set for NSS(F)., 161

 Φ_n , Markov chain value at time n, 3

 \mathcal{M} , Space of Borel probability measures, 19

 P_x , Probability conditional on $\Phi_0 = x$, 16

IR. 521

 Σ_{μ} , σ -field of P_{μ} -invariant events, 417 vec (B), 409

 $\bar{p}(M)$, upper tail of renewal sequence, 453

 $\overline{P}_k(x,\,\cdot\,)$, Cesaro average of $P^k,\,290$

 Φ , Markov chain, 3

 Φ_a Chain with transition function K_a , 122

 Φ_a , Sampled chain, 123

 $\mathcal{B}(X)$, σ -field of subsets of X, 56

 $\check{P}(x_i, A)$ The split transition function, 106

 $\check{\boldsymbol{\alpha}}$ The atom in $\mathcal{B}(\check{\mathsf{X}})$, 107

 $\dot{\Phi}$, The split chain, 106

 $C_c(X)$, continuous functions on state with compact support, 147

 $\mathcal{B}(\mathbb{R})$, 521

 $\mathcal{B}^+(X)$, Sets with $\psi(A) > 0$, 91

 $\mathcal{F}_n^{\Phi} := \sigma(\Phi_0, \dots, \Phi_n), 70$

 $\mathcal{F}_{\zeta}^{\Phi} := \{ A \in \mathcal{F} : \{ \zeta = n \} \cap A \in \mathcal{F}_{n}^{\Phi}, n \in \mathbb{Z}_{+} \}, 73$

 $\mathcal{C}(X)$, 525

 $\mathcal{C}(X)$, Continuous bounded functions on X, 132

 $C_0(X)$, continuous functions vanishing at ∞ , 527

 $\delta_x(A) = P^0(x, A)$, the Dirac measure,

 $\eta_A := \sum_{n=1}^{\infty} 1 \{ \Phi_n = A \}, 71$

 γ_q^2 , Limiting variance in the CLT, 416

 \hat{g} , Solution to the Poisson equation., 436

 $\mathbb{1}_{B}$, Indicator function of B, 69

 λ^* A split measure on bcx, 106

 μ^{Leb} Lebesgue measure, 96

 μ^{Leb} , Lebesgue measure on IR, 521

 \leftrightarrow , Communicates with, 84

 \mathbb{Z}_+ , non-negative integers, 3

 \overline{A} , Points from which A is accessible, 94

 $\overline{A}(m)$, 94

 π , Invariant measure, 234

 ψ , Maximal irreducibility measure, 90

 \rightarrow , Leads to , 84

 $\sigma_A := \min\{n \ge 0 : \Phi_n \in A\}, 71$

 $\sigma_{\alpha}(j)$, times of visits to α , 422

X, State space, 56

≻, Absolute continuity, 82

 $_{A}P^{n}(x,B) := P_{x}(\Phi_{n} \in B, \tau_{A} \geq n), 74$

 $\tau_A := \min\{n \ge 1 : \Phi_n \in A\}, 71$

 τ_A , First entry time to A, 16

 $\tau_A(1) := \tau_A, 71$

 $\tau_A(k) := \min\{n > \tau_A(k-1) : \Phi_n \in A\},$

 θ^k , The k^{th} order shift operator on Ω ,

 $\widetilde{Q}\{A\}$ An invariant random variable,

 $\tilde{\pi}\{A\}$ An invariant random variable,

 φ , Irreducibility measure, 88

 $\{\Phi \in A \text{ i.o.}\}, 204$

 $a_x(n) := P_x(\tau_{\alpha} = n), 317$

 $d(\boldsymbol{\alpha})$ The period of $\boldsymbol{\alpha}$, 118

 h_Y Almost everywhere invariant function., 417

 $m_n(t)$ Interpolation of $M_n(g)$, 440

 q_j , Probability of j arrivals in one service in the M/G/1 queue, 64

- $s_j(f)$, the sum of $f(\Phi_i)$ between visits to atom, 422
- $s_n(t)$, Interpolation of $S_n(\overline{g})$, 440
- $t_y(n) := {}_{\alpha}P^n(\boldsymbol{\alpha}, y), 317$
- $u(n) := \sum_{k=0}^{\infty} P(Z_k = n), 182$
- $u(n) := \mathsf{P}_{\alpha}(\Phi_n = \boldsymbol{\alpha}), 317$
- J, Normalized interarrival times for the Moran dam, 50, 53
- $\mathbf{W} = \{W_n\}, A \text{ disturbance process, } 24$
- **Z**, A discrete time renewal process, 45
- Φ , Markov chain, 67
- Φ^m , The m-skeleton chain, 69
- $\mu_k \xrightarrow{\text{w}} \mu_{\infty}$, weak convergence of μ_k to μ_{∞} , 147
- $\rho(F)$, maximum eigenvalue modulus, 144
- A^0 , Points from which A is inaccessible, 94
- $\operatorname{Var}(u), \operatorname{Variation}$ in a renewal sequence, 329