

Homework #3

(10 questions; 2 pages; total points = 73)

Due: In class by 6:10 p.m. **sharp** on Thursday March 30. **Warning:** Late homeworks, even by one minute, will be penalised (as discussed on the course web page).

Note: You are welcome to discuss these problems in general terms with your classmates. However, you should figure out the details of your solutions, and write up your solutions, entirely on your own. Directly copying other solutions is strictly prohibited!

[Point values are indicated in square brackets. It is very important to **EXPLAIN** all your solutions very clearly – correct answers poorly explained will **NOT** receive full marks.]

Include at the top of the first page: Your name and student number, and whether you are enrolled in STA447 or STA2006.

1. Let $\{X_n\}$ be the usual Gambler's ruin Markov chain with $p = 1/2$, $X_0 = a = 5$, and $c = 20$. Let $T = \inf\{n \geq 0 : X_n = 0 \text{ or } c\}$ as usual, and let $U = T - 1$.

(a) [2] Compute $\mathbf{E}(X_T)$.

(b) [5] Compute $\mathbf{E}(X_U)$. [Hint: If $X_T = c$, then what must X_U equal? Or, if $X_T = 0$, then what must X_U equal?]

(c) [3] Determine if $\mathbf{E}(X_T) = \mathbf{E}(X_0)$, and if $\mathbf{E}(X_U) = \mathbf{E}(X_0)$. Relate these facts to the Optional Stopping Corollary.

2. [5] Let $\{X_n\}$ be simple symmetric random walk, with $X_0 = 0$, and let $S = \inf\{n \geq 0 : X_n = -5\}$. Prove that $\lim_{M \rightarrow \infty} \mathbf{E}(X_M | S > M) = \infty$. [Hint: Let $T_M = \min(S, M)$, and apply the Optional Stopping Lemma.]

3. Let $\{X_n\}$ be a Markov chain on the state space $S = \{1, 2, 3, \dots, 100\}$, with initial probabilities given by $\nu_{30} = \nu_{40} = 1/2$, and with transition probabilities given by $p_{1,1} = p_{100,100} = 1$, $p_{99,100} = p_{99,98} = 1/2$, and for $2 \leq i \leq 98$, $p_{i,i-1} = 2/3$ and $p_{i,i+2} = 1/3$. Let $T = \inf\{n \geq 0 : X_n = 1 \text{ or } 100\}$.

(a) [3] Compute $\mathbf{P}(X_2 = 41)$.

(b) [4] Determine whether or not $\{X_n\}$ is a martingale.

(c) [4] Compute $\mathbf{P}(X_T = 1)$. [Hint: Don't forget the Optional Stopping Corollary.]

4. Suppose we repeatedly flip a fair coin.

(a) [5] Compute the expected value of the number of flips until we first see the pattern "HTHT".

(b) [5] Repeat part (a) for the pattern "HTTH".

5. [5] Let $\{B_t\}_{t \geq 0}$ be Brownian motion. Compute $\mathbf{Var}(B_5 B_8)$. [Hint: You may use without proof that if $Z \sim \text{Normal}(0, 1)$, then $\mathbf{E}(Z) = \mathbf{E}(Z^3) = 0$, $\mathbf{E}(Z^2) = 1$, and $\mathbf{E}(Z^4) = 3$.]

6. [8] Let $\{B_t\}_{t \geq 0}$ be Brownian motion. Let $\theta \in \mathbf{R}$, and let $Z_t = \exp(\theta B_t - \theta^2 t/2)$. Prove that $\{Z_t\}_{t \geq 0}$ is a martingale. [Hint: You may use without proof the fact that if $W \sim \text{Normal}(\mu, \sigma^2)$, and $a \in \mathbf{R}$, then $\mathbf{E}[e^{aW}] = e^{\mu a + \sigma^2 a^2/2} < \infty$.]

7. Let $\{B_t\}_{t \geq 0}$ be Brownian motion, let $X_t = x_0 \exp(\mu t + \sigma B_t)$ be the stock price model (where $\sigma > 0$), and let $D_t = e^{-rt} X_t$ be the discounted stock price.

(a) [4] Show that if $\mu = r - \frac{\sigma^2}{2}$, then $\{D_t\}$ is a martingale. [Hint: Don't forget the previous question.]

(b) [6] Show that if $\mu = r - \frac{\sigma^2}{2}$, then

$$\mathbf{E}\left[e^{-rS} \max(0, X_S - K)\right] = X_0 \Phi\left(\frac{(r + \frac{\sigma^2}{2})S - \log(K/X_0)}{\sigma\sqrt{S}}\right) - e^{-rS} K \Phi\left(\frac{(r - \frac{\sigma^2}{2})S - \log(K/X_0)}{\sigma\sqrt{S}}\right),$$

where $\Phi(u) = \int_{-\infty}^u \frac{1}{\sqrt{2\pi}} e^{-v^2/2} dv$ is the cdf of a standard normal distribution. [Hint: Write the expectation as an integral with respect to the density function for B_S . Then, break up the integral into the part where $X_S - K \geq 0$ and the part where $X_S - K < 0$.] [Note: This is the famous "Black-Scholes formula". You do not need to memorise it.]

8. Let $\{N(t)\}_{t \geq 0}$ be a Poisson process with intensity $\lambda = 3$.

(a) [3] Compute $\mathbf{P}[N(6) = 2 \mid N(8) = 4]$.

(b) [4] Compute $\mathbf{P}[N(6) = 2 \mid N(8) = 4, N(3) = 1]$.

(c) [4] Compute $\mathbf{E}([N(8) - N(5)][N(7) - N(2)])$. [Hint: You may use without proof the fact that if $Y \sim \text{Poisson}(m)$, then $\mathbf{E}(Y) = \mathbf{Var}(Y) = m$.]

9. Let $\{X(t)\}_{t \geq 0}$ be a continuous-time Markov process on the state space $S = \{1, 2, 3\}$. Suppose that to first order as $t \searrow 0$, its transition probabilities are given by

$$P^{(t)} = \begin{pmatrix} 1 - 7t & 7t & 0 \\ 0 & 1 - 3t & 3t \\ t & 2t & 1 - 3t \end{pmatrix} + o(t).$$

(a) [3] Compute the generator matrix G for this process.

(b) [4 bonus points] Find (with explanation) a stationary distribution $\{\pi_i\}$ for this process.

(c) [3 bonus points] Determine whether or not $\lim_{t \rightarrow \infty} p_{ij}^{(t)} = \pi_j$ for all $i, j \in S$.

10. [FOR PRACTICE ONLY, NOT TO HAND IN.] Let $\{X(t)\}_{t \geq 0}$ be a continuous-time Markov process on the state space $S = \{1, 2\}$, with generator $G = \begin{pmatrix} -3 & 3 \\ 4 & -4 \end{pmatrix}$. Compute $p_{21}^{(t)}$ for all $t > 0$ using (a) eigenvectors, and (b) differential equations.

[END; total points = 73]